arXiv:1207.6653v2 [hep-th] 2 Oct 2012

Inflation with Negative Λ

James B. Hartle,[1] S.W. Hawking,[2] and Thomas Hertog[3]

[1] *Department of Physics, University of California, Santa Barbara, 93106, USA*
[2] *DAMTP, CMS, Wilberforce Road, CB3 0WA Cambridge, UK*
[3] *Institute for Theoretical Physics, KU Leuven, 3001 Leuven, Belgium and*
International Solvay Institutes, Boulevard du Triomphe, ULB, 1050 Brussels, Belgium

The evolution of the universe is determined by its quantum state. The wave function of the universe obeys the constraints of general relativity and in particular the Wheeler-DeWitt equation. For non-zero Λ, we show that the complexified solutions of the Wheeler-DeWitt equation at large volume have two regions in which geometries are asymptotically real. In one the histories are Euclidean asymptotically anti-de Sitter, in the other they are Lorentzian asymptotically de Sitter. We illustrate this by an explicit calculation in a homogeneous, isotropic minisuperspace model with negative Λ and a scalar moving in a negative potential. It is shown that the wave function in this theory can predict an ensemble of inflationary universes that asymptote to de Sitter space.

PACS numbers: 98.80.Qc, 98.80.Bp, 98.80.Cq, 04.60.-m

Our large scale observations of the universe are of its classical behavior. The isotropic accelerated expansion and the large scale structure in the CMB and the galaxy distribution are just two examples. The laws that govern such features can be presumed to be fundamentally quantum mechanical. A quantum system behaves classically when the probabilities implied by the quantum state are high for coarse-grained histories with correlations in time governed by deterministic equations of motion. That is true whether the system is a tennis ball in flight or the evolving universe as a whole.

The form of the emergent deterministic equations of motion may be only distantly related to the equations defining the underlying quantum theory. In particular the parameters occurring in the effective equations may not be the same as those in the equations of the underlying theory. In this paper we show that in a natural formulation of quantum cosmology fundamental theories with a negative cosmological constant may lead to classical equations of motion with a positive effective cosmological constant consistent with the observed accelerated expansion [1]. A more detailed and more general exposition of this work is in [2].

Classical Prediction in Quantum Cosmology. A quantum state of a closed universe is represented by a wave function Ψ on the superspace of three-geometries and matter fields on a closed spacelike surface Σ. We consider a minisuperspace model with a negative cosmological constant Λ and a single scalar matter field ϕ with potential $V = (1/2)m^2\phi^2$. We take m^2 to be negative but within the Breitenlohner-Freedman (BF) range $m^2_{BF} < m^2 < 0$, where $m^2_{BF} = -9/(4\ell^2)$ with $1/\ell^2 \equiv -\Lambda/3$. The geometries are restricted to be spatially homogeneous, isotropic and closed. The spatial geometries on a three-sphere can then be characterized by a scale factor b and a value χ of the homogeneous field ϕ, rescaled from its usual value by a factor $\sqrt{(4\pi/3)}$ to simplify subsequent equations. Three metrics describing possible three geometries are given by

$$d\Sigma^2 = b^2 d\Omega_3^2 \tag{1}$$

A four-geometry is described by a one-parameter sequence of such scale factors. Cosmological wave functions are functions of b and χ, $\Psi = \Psi(b,\chi)$.

An ensemble of classical histories is predicted when the wave function is well approximated by a semiclassical form (or a sum of such forms)

$$\Psi(b,\chi) = A(b,\chi)\exp(iS(b,\chi)/\hbar) \tag{2}$$

where S varies rapidly compared to A. The classical histories are the integral curves of S, that is the solution to the Hamilton-Jacobi relations $p = \nabla S$ relating the momenta involving time derivatives of the variables to the derivatives of S. The probability of a classical history passing through (b,χ) is proportional to $|A(b,\chi)|^2$.

Asymptotic Structure of the Wheeler-DeWitt Equation. Cosmological wave functions must satisfy the operator forms of the constraints of general relativity. The three momentum constraints are satisfied automatically as a consequence of the symmetries of this simple model. The operator form of the remaining Hamiltonian constraint is the Wheeler-DeWitt equation (WDWE). With the simplest operator ordering this is

$$\left(-\frac{\hbar^2}{\eta^2}\frac{d^2}{db^2} + \frac{\hbar^2}{\eta^2 b^2}\frac{d^2}{d\chi^2} + b^2 + \frac{1}{\ell^2}b^4 - m^2\chi^2 b^4\right)\Psi = 0. \tag{3}$$

where $\eta \equiv 3\pi/2$. Defining $\check{I}$ by

$$\Psi(b,\chi) \equiv \exp[-\eta\check{I}(b,\chi)/\hbar] \tag{4}$$

the WDWE becomes a non-linear equation for $\check{I}$. In certain regions of superspace the terms that explicitly involve $\hbar$ may be negligible compared to those that don't. When this semiclassicality condition is satisfied

$$-\left(\frac{d\check{I}}{db}\right)^2 + \frac{1}{b^2}\left(\frac{d\check{I}}{d\chi}\right)^2 + b^2 + \left(\frac{1}{\ell^2} - m^2\chi^2\right)b^4 = 0 \tag{5}$$

which is independent of $\hbar$. Substituting a solution of (5) into (4) gives a leading order in $\hbar$ semiclassical approximation to the wave function.

The semiclassicality condition is satisfied for sufficiently large b provided $\Lambda \neq 0$. This can be established self consistently by assuming it holds, solving (5) for large b and checking that the neglected $\hbar$ terms in the WDWE are negligible [2]. Asymptotically in b sufficiently general solutions to the Hamilton-Jacobi equation (5) are defined by the expansion

$$\check{I}(b,\chi) = \frac{k_3}{\ell} b^3 + k_2 b^2 + \frac{\ell}{2} k_1 b + k_- b^{\lambda_-} + k_0 + \cdots \tag{6}$$

where the k's are all functions of χ and $\lambda_\pm \equiv (3/2)(1 \pm \sqrt{1+4\ell^2 m^2/9})$. Substituting this in (5) yields the following equations that determine the leading coefficient functions,

$$-9k_3^2 + (k_3')^2 + 1 - \ell^2 m^2 \chi^2 = 0, \tag{7a}$$

$$-6k_3 k_2 + k_3' k_2' = 0, \tag{7b}$$

$$-3k_3 k_1 + k_3' k_1' - 4k_2^2 + (k_2')^2 + 1 = 0, \tag{7c}$$

$$-3\lambda_- k_3 k_- + k_3' k_-' = 0, \tag{7d}$$

$$(2/\ell) k_0' k_3' + k_2' k_1' - 2k_2 k_1 = 0, \tag{7e}$$

a prime denoting the derivative with respect to χ.

The first three of these equations are a closed set for the first three coefficients in (6). Were there no field they would have the unique solution $k_3 = 1/3$, $k_2 = 0$, and $k_1 = 1$. We use these values as boundary conditions to fix a unique solution for the leading terms at small χ of the first three coefficients. The small χ behavior of these terms is therefore universal — the same for all wave functions satisfying the WDWE. By contrast, the large χ behavior as well as the remaining terms in (6) depend on the specific wave function.

From (6) it follows that the semiclassicality condition holds at large b as assumed, and that the general solution $\Psi(b,\chi)$ of the WDWE has the asymptotic form

$$A_+ \exp[-\eta \check{I}(b,\chi)/\hbar] + A_- \exp[+\eta \check{I}(b,\chi)/\hbar] \tag{8}$$

for arbitrary constants $A_\pm$.

Any solution of (5) with a sufficient number of arbitrary constants will determine a class of solutions to the equations of motion of general form. Specifically, sufficiently general solutions of the Hamilton-Jacobi equation (5) imply general first integrals of the Einstein equations. Hence the expansion (6) encodes the Fefferman-Graham (FG) asymptotic expansion of solutions to the Einstein equations with a negative cosmological constant [3]. The universal behavior at small χ of the first three terms in (6) corresponds to the universal terms in the FG expansion.

Two Real Domains. To find the ensemble of classical histories predicted by (8) we search for domains of asymptotic superspace that correspond to real three-geometries and field configurations where the wave function is a sum of semiclassical terms like (2), with S varying rapidly compared to A. There are two domains representing real three geometries corresponding to real values of b and purely imaginary values $b = i\hat{b}$, $\hat{b}$ real. The three-metrics in the two domains have opposite signatures, but signature is not an observable property of a metric. We take the classical ensemble to consist of the classical histories from both real domains. This is a generalization of the usual framework for quantum cosmology, but a natural one, and motivated by holography as we will shortly describe.

A Model Classical Ensemble. To exhibit the expanding histories explicitly we first restrict to the region of superspace where χ is small so the field can be treated perturbatively. The small χ solutions of (7) for the first three, universal terms in (6) are [9]:

$$k_3 = \frac{1}{3} + \frac{1}{2}\lambda_- \chi^2, \quad k_2 = 0, \quad k_1 = 1 + \frac{3}{4}\chi^2. \tag{9}$$

The leading non-universal coefficients are given by

$$k_- = K\chi, \quad k_0 = J \tag{10}$$

where K and J are complex constants (independent of χ). They are are not determined by the Hamilton-Jacobi equation but depend on the specific wave function. The relative probabilities of the predicted classical histories therefore depend on them [4].

Since k_3 in (9) is real no classical histories are predicted for the domain where b is real because there $\check{I}$ is approximately real and there is no imaginary part that varies rapidly. However, in the domain where $b = i\hat{b}$ the wave function (8) will be a sum of potentially semiclassical forms (2) with $S = \eta\check{S}$, $\check{S} = -\mathrm{Im}\check{I}$, and the leading order asymptotic behavior

$$\check{S}(\hat{b},\chi) = \frac{\hat{b}^3}{3\ell} + \frac{1}{2\ell}\lambda_- \chi^2 \hat{b}^3 + \cdots \tag{11}$$

which varies rapidly for large $\hat{b}$ so that the wave function predicts an ensemble of classical histories.

The Lorentzian histories that comprise the classical ensemble are the integral curves of $\check{S}$. The $\check{S}$ above is, in fact the action of a homogeneous and isotropic, asymptotically de Sitter history with $\Lambda = 3/\ell^2$, perturbed by a scalar field moving in a quadratic potential with positive mass $-m^2$.

The ensemble of asymptotic classical histories can be obtained explicitly by solving the Hamilton-Jacobi equations connecting the momenta to the gradients of $\check{S}$ and defining the integral curves. For this simple model these equations are

$$\frac{d\hat{b}}{dt} = \frac{1}{\hat{b}}\frac{\partial \check{S}}{\partial \hat{b}}, \quad \frac{d\chi}{dt} = -\frac{1}{\hat{b}^3}\frac{\partial \check{S}}{\partial \chi} \ . \tag{12}$$

Substituting (9) and (10) in (6) we find for the asymptotic scale factor, with $u \equiv \exp(-t/\ell)$,

$$\hat{b}(t) = \frac{2\ell}{u}[1 + u^2 - \frac{3}{4\alpha^2}u^{2\lambda_-} + \cdots] \tag{13}$$

and for the asymptotic scalar field [10]

$$\chi(t) = \alpha u^{\lambda_-} + \beta u^{\lambda_+} + \cdots \tag{14}$$

where α is an integration constant of the first order equation for χ in (12) and $\beta = -\mathrm{Im}(i^{\lambda_-}K)/(\lambda_+ - \lambda_-)$. Eqs (13) and (14) describe an ensemble of asymptotic deSitter spaces perturbed by a homogeneous decaying scalar field [5].

As ℓ approaches zero the values of b for which the asymptotic expansion (6) holds become larger and larger. If the cosmological constant vanishes then there can still be classical histories (e.g. [4]) but their existence and character depends on the specific form of the theory and the wave function. If there is no matter then there are no classical histories when $\Lambda = 0$.

Slow Roll Inflation. Beyond the perturbative regime, there is a solution of (7) which at large χ behaves as

$$k_3(\chi) \approx \frac{\ell m}{3}\chi, \quad k_2 = 0, \quad k_1 \approx 1/lm\chi \tag{15}$$

and at small χ tends to (9). This solution for k_3 is shown in Fig 1. The leading contribution to the asymptotic action in the large χ regime is then

$$\check{S}(\hat{b}, \chi) \sim \frac{m\chi\hat{b}^3}{3} + \cdots \ . \tag{16}$$

which is characteristic of cosmologies undergoing slow roll scalar field inflation [4, 6]. Indeed the leading order solutions of (12) derived from (16) exhibit the familiar slow roll behavior, $\chi(t) = \chi(0) - mt/3$.

This shows that the presence of a regime of classical slow roll inflation can be deduced from the semiclassical asymptotic form of the wave function representing a quantum state. That could be useful in a holographic study of which quantum states predict slow roll inflation.

Holography. The Euclidean AdS/CFT correspondence provides a promising route towards a more precise formulation of wave functions in gravitational theories with a negative cosmological constant [7, 8]. In particular it provides a novel way to compute the relative amplitude of different configurations in the large volume regime. It is therefore interesting to understand our results in this context.



FIG. 1: A solution of (7a) for the leading, universal coefficient function k_3 that specifies the asymptotic wave function. The approximately linear behavior for large χ is characteristic of field driven, slow roll inflation.

The duality makes a distinction between the universal contributions to the asymptotic wave function, which grow as a function of the scale factor [11], and the non-universal terms that provide a finite contribution and govern the relative weight of different configurations. The duality conjectures that the latter are related to the partition function Z of a dual Euclidean field theory defined on the conformal boundary.

In the perturbative solution of the minisuperspace model given above the universal terms are given in (9) and the non-universal terms in (10). In a regime where the low energy gravity approximation can be trusted the duality then states that, to leading order in $\hbar$ and at large volume,

$$Z[\tilde{\chi}] = \exp(-\eta(K\tilde{\chi} + J)/\hbar) \tag{17}$$

where Z is defined by

$$Z[\tilde{\chi}] \equiv \langle \exp \int d^3x \sqrt{\tilde{\gamma}}\tilde{\chi}\mathcal{O}\rangle_{QFT} \tag{18}$$

with $\mathcal{O}$ the dual operator that couples to the source $\tilde{\chi} = b^{\lambda_-}\chi$ induced by the bulk scalar. Differentiation of Z with respect to $\tilde{\chi}$ yields $\langle\mathcal{O}\rangle$ which provides an alternative way to calculate K in (17).

The bulk scalar enters as an external source in the dual field theory that turns on a deformation. In field theory applications of AdS/CFT the value of $\tilde{\chi}$ is usually held fixed. By contrast, in a cosmological context one is interested in the wave function of the universe as a function of χ; the dependence of Z on the value of $\tilde{\chi}$ yields a measure on different configurations on Σ [8].

Hence in cosmology one is led to consider all deformations induced by a given quantum state that correspond to real boundary configurations and for which the partition function converges. In the simple model discussed here this means one should consider purely real deformations, corresponding to real values of b, as well as deformations for which $\tilde{\chi}$ has the phase $\exp[i\lambda_-\pi/2]$.

Together these generate the two real domains of asymptotic superspace discussed earlier.

In a dual formulation of the wave function, the difference between both domains thus shows up in the phase of the scalar deformation. Since the phase of $\tilde{\chi}$ in general has an effect on the partition function this means that the asymptotic wave function will be a slightly different function of χ in both domains. We illustrated this with the example of the no-boundary wave function in [2].

Discussion. Accelerated expansion in a theory with a negative cosmological constant is surprising from the perspective of theories which posit a fixed background spacetime as a classical arena for quantum physics. But in a quantum universe there is no general reason to assume that some degrees of freedom behave classically and others quantum mechanically. Instead both backgrounds and fluctuations should be treated quantum mechanically, as we do here. In a fully quantum theory of cosmology the probabilities for histories following from a quantum state determine what behaves classically and what does not. Furthermore the form of the resulting classical equations is given by the quantum state and not by assumption.

This paper has used a natural framework for the prediction of probabilities of classical histories to show (in a simple model) that accelerated expansion can be a low energy prediction of quantum theories with a negative cosmological constant. Specifically we have shown that, in models with a non-zero cosmological constant, wave functions satisfying the constraints of general relativity have a universal semiclassical asymptotic structure for large spatial volumes. This asymptotic structure implies that a wave function in a gravitational theory with a negative cosmological constant describes both a set of asymptotically Euclidean AdS histories as well as an ensemble of histories which expand, driven by a positive effective cosmological constant. The relative probabilities of different histories depend on the specific wave function. The generalization of these results to include inhomogeneities is described in [2].

Our results do not arise from analytically continuing the parameters of a background from one theory to another. Rather we have one theory which predicts an ensemble of possible backgrounds, not just one. We exploit the analytic properties of wave functions that solve the WDWE equation to generalize the process of classical prediction in one theory with a single set of parameters.

Prospects. One may ask how the models discussed in this paper can be embedded in fundamental theory. Examples of models to which our results apply are given by the consistent truncations of the low energy limit of M theory compactified on S^7 involving only AdS gravity and one or several scalars with negative mass $m^2 = -2/\ell^2$. The scalar potential in these truncations acts as a positive effective potential in the 'cosmological' domain of the theory, and the WDWE has solutions that predict an ensemble of classical universes with an epoch of scalar field driven inflation.

However, the predicted inflationary histories are likely to be unstable when viewed as low-energy solutions in the full theory, especially towards the end of the slow roll regime. This is because this also contains scalars with a positive potential around the AdS vacuum. These give rise to instabilities in the cosmological region where they have negative effective mass. The requirement of a (meta)stable de Sitter era at late times may thus act as a vacuum selection principle that excludes this class of compactifications and picks out a different set.

Alternatively one might speculate that the above instability, which appears to be rather generic in Freund-Rubin compactifications, can be incorporated in a theory of reheating in a scenario akin to hybrid inflation, where one vacuum decays at the end of inflation to a vacuum with a lower cosmological constant. A complete description of cosmology would then require a model which describes both the unstable early phase of accelerated expansion and the stable de Sitter phase at late times.

Acknowledgments: JH and SWH thank Marc Henneaux and the International Solvay Institutes for their hospitality. The work of JH was supported in part by the US NSF grant PHY08-55415. The work of TH was supported in part by the FWO-Vlaanderen under the Odysseus program.

---

[1] **WMAP** Collaboration, E. Komatsu *et al.*, Astrophys. J. Suppl. **192**, 18 (2011), arXiv:1001.4538

[2] J.B. Hartle, S.W. Hawking, T. Hertog, arXiv:1205.3807.

[3] C. Fefferman, C. R. Graham in *Elie Cartan et les Mathématiques d'aujourd'hui* (Astérisque, 1985) 95.

[4] J.B. Hartle, S.W. Hawking, T. Hertog, Phys. Rev. D **77**, 123537 (2008), arXiv:0803.1663; Phys. Rev. Lett. **100**, 202301 (2008), arXiv:0711.4630.

[5] A. Starobinsky, Pis'ma Zh. Eksp. Teor. Fiz. **37**, 55 (1983) [JETP Lett. **37**, 66 (1983)]

[6] G.W. Lyons, *Complex solutions for the scalar field model of the universe,* Phys. Rev. D **46**, 1546-1550 (1992).

[7] G. T. Horowitz, J. M. Maldacena, JHEP **0402**, 008 (2004), hep-th/0310281

[8] T. Hertog and J.B. Hartle, JHEP **1205**, 095 (2012), arXiv:1111.6090

[9] Replacing λ_- by λ_+ in the first equation of (9) also gives a perturbative solution but a numerical integration of the non-linear eq (7a) shows that (9) is the generic one.

[10] When $\lambda_- < 1/2$ there is an additional universal term $(-3\alpha/8)u^{2+\lambda_-}$.

[11] In field theory applications of AdS/CFT these are treated as UV divergences and cancelled by counterterms.